Feral Swine Information

Feral swine pose significant health threats to domestic pigs and even to people. These wild hogs are appearing in Georgia in areas that did not have populations in the past. There is evidence of increasing transportation and relocation for hunting purposes.

Feral swine brought into Georgia from another State for any purpose other than slaughter are required to have an entry permit from the office of the Georgia State Veterinarian and a certificate of veterinary inspection with negative results for brucellosis and pseudorabies within the past 30 days. Once inside the State, feral swine and domestic swine must have the same testing for these diseases at every change of ownership.

Testing of feral swine in Georgia from 1991 through January 1, 1999, reveals the following:

Hunter-Killed or Confined Feral Swine

- Total tested 1.801
- Total positive for pseudorabies 507, or 28.2 percent
- Total positive for swine brucellosis 73, or 4.1 percent

Swine brucellosis and pseudorabies are both transmissible to domestic pigs, and swine brucellosis can be transmitted to people. Everyone in contact with feral swine should wear rubber gloves.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, race, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720–2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326–W, Whitten Building, 1400 and Independence Avenue, SW, Washington, DC 20250–9410 or call (202) 720–5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Photo credit: The image on the cover was taken by APHIS veterinarians Mike Duffy and David Larimer, in Toombs County, GA, especially for this publication. The new image of a feral pig inside the leaflet was supplied by the Southeastern Cooperative Wildlife Disease Study in Athens, GA.

Issued July 1997 Slightly revised September 2000

Keep Georgia's Swine Herd Safe

Do not relocate feral swine without a negative test for pseudorabies and brucellosis.

Do not introduce feral swine to a hunting area without first obtaining the landowner's permission.

Do not release feral swine into or near a domestic swine operation.

Do not handle feral swine without wearing rubber gloves.



For More Information

USDA's Animal and Plant Health Inspection Service (APHIS) has published a color brochure on feral swine. If you'd like a free copy of Wild Pigs: Hidden Danger for Farmers and Hunters (Agriculture Information Bulletin No. 620), write to

Management Services Division USDA, APHIS 4700 River Road, Unit 1 Riverdale, MD 20737–1229

If you have any questions regarding feral swine, please contact

Dr. Carter Black Georgia Department of Agriculture Atlanta, GA 1–800–282–5852, ext. 3667 or 404–656–3667

Area Epidemiologist USDA, APHIS, Veterinary Services Conyers, GA 770–922–7860 Voice mail ext. 107 USDA, APHIS, Veterinary Services 1498 Klondike Road, Suite 200 Conyers, GA 30094 770–922–7860

And Cooperating

Georgia Department of Agriculture Agriculture Building Capitol Square Atlanta, GA 30334 404–656–3671



U.S. Department of AgricultureAnimal and Plant Health Inspection Service

Georgia Department of Agriculture

APHIS 91-55-036

Feral Swine Information

